



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

allied in its species to Eastern Asia, a fact brought out not only by a similarity in common species or large genera, but in very many cases by peculiar species and genera represented by two species only, one in Eastern America, the other in Eastern Asia. In Western America there is no such resemblance, making a greater difference between the floras of the eastern and western parts of the United States than between Eastern America and Eastern Asia. In the west there is a large commingling of Mexican or Southern species that have crept northward, guided by the mountain ranges. Hence the conclusion is arrived at that the similarity between the species of Eastern America and Eastern Asia is explained by the continuity of the continents to the north in the Cretaceous or Miocene and a consequent commingling of species, and that the Glacial period drove these species southward along the two continents farther even than they appear now. With the retreat of the glacier and the return of a milder climate these plants would creep northward again, but the enormous height of the Rocky Mountain and Sierra Nevada regions would retain the glaciers there long after they had disappeared from the eastern half of the continent. When eventually these alpine glaciers did succumb to a milder climate, the old Asiatico-American species having either established themselves elsewhere or been driven into the sea towards the south, the field would be clear for the advance of the Mexican forms and we find them even up in British America. Hence to state it all in one sentence, our Eastern flora has come from the North and our Western flora from the South.—J. M. C.

CAUCALIS ANTHERISCUS.—I have found thoroughly naturalized in the woods back of Cincinnati and remote from dwellings, *Caucalis Antheriscus*. The name was kindly furnished by J. W. Congdon. Prof. Watson, to whom I sent a specimen, informs me that the plant was found in 1872 by Judge G. W. Clinton, near Buffalo, N. Y. It is firmly established in this locality and promises, I think, to become a troublesome weed.—C. G. LLOYD.

NOTE ON PANICUM LITTORALE, VASEY, BY GEN. WM. MUNRO.—PANICUM LITTORALE, Vasey, is undoubtedly *P. repens*, L.,—one specimen the dwarfish form which he described in the second edition of Sp. Plant. p. 87. It is absolutely identical with specimens which I have before me in DeCandolle's herbarium from Crete and Gibraltar, where I have also collected it myself. It is also *P. arenarium*, Brotero.

There has always been considerable confusion about *P. repens*, L. The *P. repens* of Burman, Fl. Ind., p. 26, tab. 11, fig. 1 (1768), where